

# 2007 NIH Asthma Guidelines Summary

## **Summary of the *Expert Panel Report 3:* *Guidelines for the Diagnosis and Management of Asthma***

National Institutes of Health (NIH)  
National Heart, Lung, and Blood Institute  
National Asthma Education and Prevention Program (NAEPP)

This information is abstracted from the 2007 NAEPP *Expert Panel Report 3: Guidelines for the Diagnosis and Management of Asthma*. To access the complete report, go to [www.nhlbi.nih.gov/guidelines/asthma/asthgdln.pdf](http://www.nhlbi.nih.gov/guidelines/asthma/asthgdln.pdf).

# Goals of Asthma Control<sup>1</sup>

## ▶ Reduce impairment

- Prevent chronic and troublesome symptoms
- Require infrequent use of inhaled short-acting beta<sub>2</sub>-agonists (SABA) for quick relief of symptoms ( $\leq 2$  days per week)
- Maintain (near) “normal” pulmonary function
- Maintain normal activity levels (including exercise and other physical activity and attendance at work or school)
- Meet patients’ and families’ expectations of and satisfaction with asthma care

## ▶ Reduce risk

- Prevent recurrent exacerbations and minimize the need for emergency department (ED) visits/hospitalizations
- Prevent progressive loss of lung function; for children, prevent reduced lung growth
- Provide optimal pharmacotherapy with minimal or no adverse effects

## Guidelines include separate but related concepts of severity, control, and responsiveness to treatment

### ▶ Severity

- The intrinsic intensity of the disease process
- Measured most easily and directly in a patient not receiving long-term control therapy
- Assessed to guide clinical decisions on appropriate medications and interventions

### ▶ Control

- The degree to which the manifestations of asthma (symptoms, functional impairments, and risks of untoward events) are minimized and the goals of therapy are met
- Guide decisions to maintain or adjust therapy

### ▶ Responsiveness

- The ease with which asthma control is achieved by therapy

## Assess asthma severity to initiate therapy

- During a patient’s initial presentation, if the patient is not currently taking long-term control medication, assess asthma severity to guide clinical decisions for initiating the appropriate medication and other therapeutic interventions

## Assess asthma control to monitor and adjust therapy

- Once therapy is initiated, the emphasis for clinical management is changed to the assessment of asthma control. Use the level of asthma control to guide decisions either to maintain or to adjust therapy

# Patients ≥12 Years of Age

## Assessing Severity: Patients ≥12 Years of Age

| Classifying Asthma Severity and Initiating Treatment in Patients ≥12 Years of Age <sup>1</sup>                      |   |  |   |   |   |
|---|---|--|---|---|---|
| Assessing severity and initiating treatment for patients who are not currently taking long-term control medications |   |  |   |   |   |
| Components of Severity  |   | Classification of Asthma Severity (≥12 Years of Age)   |   |   |   |
|   |   | Intermittent   | Persistent  |   |   |
|   |   |  | Mild  | Moderate  | Severe  |
| IMPAIRMENT*   | Symptoms  | ≤2 days/week   | >2 days/week but not daily  | Daily   | Throughout the day  |
|   | Nighttime awakenings  | ≤2x/month  | 3-4x/month  | >1x/week but not nightly  | Often 7x/week   |
|   | Short-acting beta <sub>2</sub> -agonist use for symptom control (not prevention of EIB) | ≤2 days/week   | >2 days/week but not daily, and not more than 1x on any day   | Daily   | Several times per day   |
|   | Interference with normal activity   | None   | Minor limitation  | Some limitation   | Extremely limited   |
|   | Lung function   | <ul style="list-style-type: none"> <li>• Normal FEV<sub>1</sub> between exacerbations</li> <li>• FEV<sub>1</sub> &gt;80% predicted</li> <li>• FEV<sub>1</sub>/FVC normal</li> </ul>  | <ul style="list-style-type: none"> <li>• FEV<sub>1</sub> &gt;80% predicted</li> <li>• FEV<sub>1</sub>/FVC normal</li> </ul> | <ul style="list-style-type: none"> <li>• FEV<sub>1</sub> &gt;60% but &lt;80% predicted</li> <li>• FEV<sub>1</sub>/FVC reduced 5%</li> </ul> | <ul style="list-style-type: none"> <li>• FEV<sub>1</sub> &lt;60% predicted</li> <li>• FEV<sub>1</sub>/FVC reduced &gt;5%</li> </ul> |
| RISK  | Exacerbations requiring oral systemic corticosteroids                                   | 0-1 per year   | ≥2 per year   |   |   |
|   |   | Consider severity and interval since last exacerbation.<br>← Frequency and severity may fluctuate over time for patients in any severity category. →<br>Relative annual risk of exacerbations may be related to FEV <sub>1</sub> . |   |   |   |
| Recommended Step for Initiating Therapy   |   | Step 1   | Step 2  | Step 3  | Step 4 or 5   |
| In 2-6 weeks, evaluate level of asthma control that is achieved, and adjust therapy accordingly.                    |   |  |   |   |   |

\*Normal FEV<sub>1</sub>/FVC: 8-19 yr: 85% | 20-39 yr: 80% | 40-59 yr: 75% | 60-80 yr: 70%

## Assessing Control: Patients ≥12 Years of Age


| Assessing Asthma Control and Adjusting Therapy in Patients ≥12 Years of Age <sup>1</sup> |   |  |   |  |
|--|---|--|---|--|
| Components of Control  |   | Classification of Asthma Control (≥12 Years of Age)  |   |  |
|  |   | Well Controlled  | Not Well Controlled   | Very Poorly Controlled   |
| IMPAIRMENT   | Symptoms  | ≤2 days/week   | >2 days/week  | Throughout the day   |
|  | Nighttime awakenings  | ≤2x/month  | 1-3x/week   | ≥4x/week   |
|  | Interference with normal activity   | None   | Some limitation   | Extremely limited  |
|  | Short-acting beta <sub>2</sub> -agonist use for symptom control (not prevention of EIB) | ≤2 days/week   | >2 days/week  | Several times per day  |
|  | FEV <sub>1</sub> or peak flow   | >80% predicted/personal best   | 60%-80% predicted/personal best   | <60% predicted/personal best   |
|  | Validated questionnaires<br>ATAQ<br>ACQ<br>ACT‡   | 0<br>≤0.75†<br>≥20   | 1-2<br>≥1.5<br>16-19  | 3-4<br>N/A<br>≤15  |
| RISK   | Exacerbations requiring oral systemic corticosteroids                                   | 0-1 per year   | ≥2 per year   |  |
|  | Progressive loss of lung function   | Evaluation requires long-term follow-up care.  |   |  |
|  | Treatment-related adverse effects   | Medication side effects can vary in intensity from none to very troublesome and worrisome. The level of intensity does not correlate to specific levels of control but should be considered in the overall assessment of risk. |   |  |
| Recommended Action for Treatment   |   | <ul style="list-style-type: none"> <li>• Maintain current step.</li> <li>• Regular follow-ups every 1-6 months to maintain control.</li> <li>• Consider step down if well controlled for at least 3 months.</li> </ul>         | <ul style="list-style-type: none"> <li>• Step up 1 step and</li> <li>• Reevaluate in 2-6 weeks.</li> <li>• For side effects, consider alternative treatment options.</li> </ul> | <ul style="list-style-type: none"> <li>• Consider short course of oral systemic corticosteroids,</li> <li>• Step up 1-2 steps, and</li> <li>• Reevaluate in 2 weeks.</li> <li>• For side effects, consider alternative treatment options.</li> </ul> |

†ACQ values are indeterminate regarding well-controlled asthma.

‡ACT = Asthma Control Test. Asthma Control Test is a trademark of QualityMetric Incorporated.

# Patients ≥12 Years of Age

## Assessing Treatment Options: Patients ≥12 Years of Age



## Patients 5-11 Years of Age


### Assessing Severity: Children 5-11 Years of Age

| Classifying Asthma Severity and Initiating Treatment in Children 5-11 Years of Age <sup>1</sup>                   |   |  |  |  |  |
|---|---|--|--|--|--|
| Assessing severity and initiating treatment in children who are not currently taking long-term control medication |   |  |  |  |  |
| Components of Severity  |   | Classification of Asthma Severity (5-11 Years of Age)  |  |  |  |
|   |   | Intermittent   | Persistent   |  |  |
|   |   |  | Mild   | Moderate   | Severe   |
| IMPAIRMENT  | Symptoms  | ≤2 days/week   | >2 days/week but not daily   | Daily  | Throughout the day   |
|   | Nighttime awakenings  | ≤2x/month  | 3-4x/month   | >1x/week but not nightly   | Often 7x/week  |
|   | Short-acting beta <sub>2</sub> -agonist use for symptom control (not prevention of EIB) | ≤2 days/week   | >2 days/week but not daily   | Daily  | Several times per day  |
|   | Interference with normal activity   | None   | Minor limitation   | Some limitation  | Extremely limited  |
|   | Lung function   | <ul style="list-style-type: none"> <li>• Normal FEV<sub>1</sub> between exacerbations</li> <li>• FEV<sub>1</sub> &gt;80% predicted</li> <li>• FEV<sub>1</sub>/FVC &gt;85%</li> </ul> | <ul style="list-style-type: none"> <li>• FEV<sub>1</sub> = &gt;80% predicted</li> <li>• FEV<sub>1</sub>/FVC &gt;80%</li> </ul> | <ul style="list-style-type: none"> <li>• FEV<sub>1</sub> = 60%-80% predicted</li> <li>• FEV<sub>1</sub>/FVC = 75%-80%</li> </ul> | <ul style="list-style-type: none"> <li>• FEV<sub>1</sub> &lt;60% predicted</li> <li>• FEV<sub>1</sub>/FVC &lt;75%</li> </ul> |
| RISK  | Exacerbations requiring oral systemic corticosteroids                                   | 0-1 per year   | ≥2 per year  |  |  |
|   |   | Consider severity and interval since last exacerbation.<br>Frequency and severity may fluctuate over time for patients in any severity category.                                     |  |  |  |
|   |   | Relative annual risk of exacerbations may be related to FEV <sub>1</sub> .   |  |  |  |
| Recommended Step for Initiating Therapy   |   | Step 1   | Step 2   | Step 3, medium-dose ICS option   | Step 3, medium-dose ICS option, or Step 4 and consider short course of oral systemic corticosteroids                         |
| In 2-6 weeks, evaluate level of asthma control that is achieved, and adjust therapy accordingly.                  |   |  |  |  |  |

### Assessing Control: Children 5-11 Years of Age

| Assessing Asthma Control and Adjusting Therapy in Children 5-11 Years of Age <sup>1</sup> |   |  |  |   |  |
|---|---|--|--|---|--|
| Components of Control   |   | Classification of Asthma Control (5-11 Years of Age)   |  |   |  |
|   |   | Well Controlled  | Not Well Controlled  | Very Poorly Controlled  |  |
| IMPAIRMENT  | Symptoms  | ≤2 days/week but not more than once on each day  | >2 days/week or multiple times on ≤2 days/week   | Throughout the day  |  |
|   | Nighttime awakenings  | ≤1x/month  | ≥2x/month  | ≥2x/week  |  |
|   | Interference with normal activity   | None   | Some limitation  | Extremely limited   |  |
|   | Short-acting beta <sub>2</sub> -agonist use for symptom control (not prevention of EIB)                                       | ≤2 days/week   | >2 days/week   | Several times per day   |  |
|   | Lung function <ul style="list-style-type: none"> <li>• FEV<sub>1</sub> or peak flow</li> <li>• FEV<sub>1</sub>/FVC</li> </ul> | >80% predicted/personal best >80%  | 60%-80% predicted personal best 75%-80%  | <60% predicted/personal best <75% predicted   |  |
| RISK  | Exacerbations requiring oral systemic corticosteroids   | 0-1 per year   | ≥2 per year  |   |  |
|   |   | Consider severity and interval since last exacerbation.  |  |   |  |
|   |   | Reduction in lung growth Evaluation requires long-term follow-up care.   |  |   |  |
| Treatment-related adverse effects   |   | Medication side effects can vary in intensity from none to very troublesome and worrisome. The level of intensity does not correlate to specific levels of control but should be considered in the overall assessment of risk. |  |   |  |
| Recommended Action for Treatment  |   | <ul style="list-style-type: none"> <li>• Maintain current step.</li> <li>• Regular follow-up every 1-6 months.</li> <li>• Consider step down if well controlled for at least 3 months.</li> </ul>                              | <ul style="list-style-type: none"> <li>• Step up 1 step and Reevaluate in 2-6 weeks.</li> <li>• For side effects, consider alternative treatment options.</li> </ul> | <ul style="list-style-type: none"> <li>• Consider short course of oral systemic corticosteroids,</li> <li>• Step up 1-2 steps, and Reevaluate in 2 weeks.</li> <li>• For side effects, consider alternative treatment options.</li> </ul> |  |

## Assessing Treatment Options: Children 5-11 Years of Age



# Patients 0-4 Years of Age

## Assessing Severity: Children 0-4 Years of Age


| Classifying Asthma Severity and Initiating Treatment in Children 0-4 Years of Age <sup>1</sup>   |   |  |   |   |                       |
|--|---|--|---|---|-----------------------|
| Assessing severity and initiating treatment in children who are not currently taking long-term control medications   |   |  |   |   |                       |
| Components of Severity   |   | Classification of Asthma Severity (0-4 Years of Age)   |   |   |                       |
|  |   | Intermittent   | Persistent  |   |                       |
|  |   |  | Mild  | Moderate  | Severe                |
| IMPAIRMENT   | Symptoms  | ≤2 days/week   | >2 days/week but not daily  | Daily   | Throughout the day    |
|  | Nighttime awakenings  | 0  | 1-2x/month  | 3-4x/month  | >1x/week              |
|  | Short-acting beta <sub>2</sub> -agonist use for symptom control (not prevention of EIB) | ≤2 days/week   | >2 days/week but not daily  | Daily   | Several times per day |
|  | Interference with normal activity   | None   | Minor limitation  | Some limitation   | Extremely limited     |
| RISK   | Exacerbations requiring oral systemic corticosteroids                                   | 0-1 per year   | ≥2 exacerbations in 6 months requiring oral systemic corticosteroids, or ≥4 wheezing episodes/1 year lasting >1 day AND risk factors for persistent asthma. |   |                       |
|  |   | ← Consider severity and interval since last exacerbation. Frequency and severity may fluctuate over time. →<br>Exacerbations of any severity may occur in patients in any severity category. |   |   |                       |
| Recommended Step for Initiating Therapy  |   | Step 1   | Step 2  | Step 3 and consider short course of oral systemic corticosteroids |                       |
| In 2-6 weeks, depending on severity, evaluate level of asthma control that is achieved. If no clear benefit is observed in 4-6 weeks, consider adjusting therapy or alternative diagnoses. |   |  |   |   |                       |

## Assessing Control: Children 0-4 Years of Age

| Assessing Asthma Control and Adjusting Therapy in Children 0-4 Years of Age <sup>1</sup> |   |  |  |   |
|--|---|--|--|---|
| Components of Control  |   | Classification of Asthma Control (0-4 Years of Age)  |  |   |
|  |   | Well Controlled  | Not Well Controlled  | Very Poorly Controlled  |
| IMPAIRMENT   | Symptoms  | ≤2 days/week   | >2 days/week   | Throughout the day  |
|  | Nighttime awakenings  | ≤1x/month  | >1x/month  | >1x/week  |
|  | Interference with normal activity   | None   | Some limitation  | Extremely limited   |
|  | Short-acting beta <sub>2</sub> -agonist use for symptom control (not prevention of EIB) | ≤2 days/week   | >2 days/week   | Several times per day   |
| RISK   | Exacerbations requiring oral systemic corticosteroids                                   | 0-1 per year   | 2-3 per year   | >3 per year   |
|  | Treatment-related adverse effects   | Medication side effects can vary in intensity from none to very troublesome and worrisome. The level of intensity does not correlate to specific levels of control but should be considered in the overall assessment of risk. |  |   |
| Recommended Action for Treatment   |   | <ul style="list-style-type: none"> <li>• Maintain current treatment.</li> <li>• Regular follow-up every 1-6 months.</li> <li>• Consider step down if well controlled for at least 3 months.</li> </ul>                         | <ul style="list-style-type: none"> <li>• Step up 1 step and Reevaluate in 2-6 weeks.</li> <li>• If no clear benefit in 4-6 weeks, consider alternative diagnoses or adjusting therapy.</li> <li>• For side effects, consider alternative treatment options.</li> </ul> | <ul style="list-style-type: none"> <li>• Consider short course of oral systemic corticosteroids,</li> <li>• Step up 1-2 steps, and Reevaluate in 2 weeks.</li> <li>• If no clear benefit in 4-6 weeks, consider alternative diagnoses or adjusting therapy.</li> <li>• For side effects, consider alternative treatment options.</li> </ul> |

# Patients 0-4 Years of Age (continued)

## Assessing Treatment Options: Children 0-4 Years of Age



**Reference:** 1. National Heart, Lung, and Blood Institute. Expert panel report 3: guidelines for the diagnosis and management of asthma: full report 2007. NIH publication 08-4051. <http://www.nhlbi.nih.gov/guidelines/asthma/asthgdln.htm>. Accessed February 28, 2008.